Les Houches Workshop: TeV Scale Physics*

The future…

Where are we? Why the TeV scale?

What physics should we be doing?

Sally Dawson
BNL
June 6, 2003

* NOT a summary….my personal views

Scaling the Mountains…..

THANKS TO THE ORGANIZERS!



Broad definition of particle physics

The Challenge: Connecting the Energy Scales*



* US version is vertical

Much Progress last year

- Many simulations reported here
- Many new tools
  - For Monte Carlos
  - For PDFs (uncertainties!)
  - For higher order corrections
  - For SUSY spectrum
  - For Higgs Properties
  - ……
- Many new ideas

Machines for Collider Physics

Tevatron



~ 9-15 fb^{-1} by Sept. 2009

HERA

Linear Collider
20??

10 fb^{-1} by 2007-2008

$\rightarrow$100 fb^{-1}

$\rightarrow$300 fb^{-1}

Looking towards 2007

W mass	**20 –30 MeV**	**LEP + Tevatron**
Top mass	**2 GeV**	**Tevatron**
Structure functions:*	**Few percent up to $Q^2 = 10^4$ GeV^2**	**HERA**
Higgs, SUSY	**Possible signal**	**Tevatron**

* Gluon PDFs critical at LHC

Physics landscape changes in 2007

Event Rates:

Process	10 fb^{-1} at LHC	Previous experiments
$W \rightarrow e\nu$	10^8	10^7 Tevatron
$Z \rightarrow e^+e^-$	10^7	10^7 LEP
$t\bar{t}$	10^7	10^4 Tevatron
$b\bar{b}$	10^{12}-10^{13}	10^9 B factories
Higgs, M_h=130 GeV	10^5	?
$\tilde{g}\tilde{g}$, m=1 TeV	10^4	--

Experimental successes of past decade put us on firm footing

We have a model

And it works to the 1% level

Gives us confidence to predict the future!



We've seen one example of gauge unification

HERA



Charged and neutral currents unify at 100 GeV

Model requires Higgs boson or something like it for consistency!

Why the TeV Scale?
We expect new physics on very general grounds

Either a light Higgs, or strong WW Scattering





Standard Model inconsistent without Higgs unless new physics around 1.3 TeV

Kolda & Murayama, hep-ph/0003170

Is Mass due to a Higgs Boson?

Precision EW Measurements:

$M_h < 211$ GeV





Note: Poor quality of fit

The Tevatron will point the way….



Increasing M_t increases M_h



Theory errors

D0 preliminary:

M_t=180.1±3.6±4.0 GeV

If there is a light SM Higgs, we'll find it







Conclusions "basically" confirmed by re-analysis

Progress in Vector Boson Fusion Studies

Note: 10 fb^{-1}

M_H= 115 GeV
combined significance ~ 5σ

Promising channel for measuring couplings



Coupling Constant Measurements

Ratios of coupling constants, quite precisely



Duehrssen

ATLAS



Coupling constants less precisely

Coupling Constant Measurements at a Linear Collider



L=500 fb^{-1}, $\sqrt{s}$=350 GeV

Battaglia & Desch, hep-ph/0101165

Global fit to Higgs Couplings:

$\delta g_h/g_h$	
M_h=120 GeV	
bb	2.1%
$\tau^+\tau^-$	3.2%
W^+W^-	1.2%
cc	3.2%

Largest error is theory!
(mostly from m_b)

Linear Collider is the place!

Can we reconstruct the Higgs potential?

$$V = \frac{M_h^{\,2}}{2} h^2 + \lambda_3 v h^3 + \frac{\lambda_4}{4} h^4$$

$$+ \sum_n C_n \frac{(h^2 - v^2)^n}{\Lambda^{(2n-4)}}$$

Fundamental test of model!



Reconstructing the Higgs potential





- λ_3 requires 2 Higgs production
- Depressing story
- No prospects for λ_4
- Hard at Linear collider too!

Can determine whether λ_3=0



Baur, Plehn & Rainwater, hep-ph/0304015

What do Higgs couplings mean?
Can they tell us about new physics?



"At this point we notice that this equation is beautifully simplified if we assume that space–time has 92 dimensions."

Precision measurements versus direct observation of new particles



New Physics Searches require
Precision Calculations

- Is it new physics?
  - Extra dimensions
  - Little Higgs
  - SUSY
  - Something really new*…..
- Or is it QCD?

*Lots of theoretical progress



Of course….we all hope for physics beyond the Standard Model

- MSSM most studied variant of SM
- Motivated by coupling unification; Higgs mass renormalization
- Definite predictions



Doesn't happen in SM

$M_A \to \infty$, light Higgs looks like SM

Focus on finding other Higgs bosons (signal for new physics beyond SM)

New decay channel H/A $\to \mu^+\mu^-$ covers much of region not excluded by LEP

New signature from $gb \to tH^+$, $H^+ \to \tau\upsilon$

H/A $\to \tau^+\tau^-$ important for intermediate tan β



Emphasis on new signatures

One example:

- Cross section for pair production of squarks, gluinos large at LHC
- Look for complicated cascade decays
- Fills difficult low M_A, tan β region

$$\tilde{g}, \tilde{q} \rightarrow ..... \rightarrow \tilde{\chi}^{\pm}, \tilde{\chi}^{0} + h$$



Datta, Djouadi, Guchait, Moortgat,

hep-ph/0303095

What does it all mean?

- If we measure SUSY masses at LHC, can we determine underlying SUSY theory?
- Extrapolate measured values with 300 fb^{-1} to M_{GUT}

Les Houches Accord





Allanach, Kraml, Porod, hep-ph0302102

We believe we understand parameters running with mass scales…..

Necessary ingredient for SUSY extrapolations

Running $\sin^2\theta_W^{eff}$



Czarnecki & Marciano

Moller Scattering,

Pure electroweak



$\sin 2\theta_W^{eff}(M_Z)$

Models with Extra Dimensions provide another good comparison point for SM

- Extra-D Models have towers of new Kaluza Klein Gravitons
- Graviton emission can measure the number of hidden dimensions
- Graviton exchange affects precision measurements,

  Drell-Yan production, missing E_T measurements….



Large Extra Dimensions change angular, mass distributions of lepton pairs

RunII search for high mass di-leptons

Sensitive to Z' and Randall-Sundrum Graviton

No excess observed





Note critical importance of understanding QCD backgrounds

First NNLO distribution: Drell Yan rapidity



Anastasiou,Dixon, Melnikov, Petriello (preliminary)

NLO, NNLO corrections significant at y=0

Calculation possible because of new techniques!

Nothing like direct production…

Drell-Yan production of KK Gravitons

in Randall-Sundrum Model

$pp \rightarrow$ Graviton KK $\rightarrow l^+l^-$





Davoudiasl, Hewett & Rizzo, hep-ph/0006041

Once we find something, we want to know what it is….

Is it a Higgs or is it a radion?

- Randall-Sundrum model of extra dimensions has radion: ϕ
- ϕ couples like Higgs, but with strength ϕ/TeV instead of H/v
- Higgs-ϕ mixing suppresses standard channel, $gg \rightarrow h \rightarrow \gamma\gamma$
- How do you know it's a radion, not a SUSY Higgs?

(Have to find both Higgs & radion)



Battaglia, DeCurtis, DeRoeck, Dominici,& Gunion, hep-ph/030425

Dominici, Grzadkowski, Gunion, Toharia, hep-ph/0206192

Hewett & Spiropulu, hep-ph/0205100

Need NLO, NNLO Results

bb→h*



Note strong μ dependence at NLO

Only relevant in SUSY with large tan β

* When is the b quark a parton????

Reduced theoretical error from reduced μ dependence



NLO: Maltoni, Sullivan, & Willenbrock, hep-ph/0301033

NNLO: Harlander & Kilgore, hep-ph/0304035

NNLO, gg→h



Bands show $.5M_h < \mu < 2\ M_h$

LO and NLO μ dependence bands don't overlap.

Need NNLO!

Resummation ↑ σ by 6% wrt NNLO [Grazzini]

NLO&NNLO results allow improved estimates of theoretical uncertainties

Harlander & Kilgore, hep-ph/0201202

Catani, deFlorian, Grazzini, hep-ph/0102227

Anastasiou & Melnikov, hep-ph/0207004

Need backgrounds to Higher Order

- Background to $h \to \gamma\gamma$:
- $qq \to \gamma\gamma$
- $gg \to \gamma\gamma$



"NLO"



Bern, de Freitas, Schmidt, hep-ph/0206194

Previous estimates over-estimated $\gamma\gamma$ background

New ATLAS analysis of $H \to \gamma\gamma$ with increased significance

More on higher order QCD

Higgs p_T spectrum at NLO QCD:



NLO corrections to shape small, 5% at y=0

Suggests using LO Monte Carlo, weighted by NNLO cross section

Need case by case study

Anastasiou, Dixon, &Melnikov, hep-ph/0211141

Glosser & Schmidt, hep-ph/0209248

Ravindran, Smith, & van Neervan, hep-ph/021114

deFlorian, Grazzini, & Kunszt, hep-ph/9902483

Progress in estimating theoretical errors

- CTEQ6 PDFs have errors
- NLO, NNLO have reduce scale dependence

Theoretical errors critical for extracting tan β….

PDF uncertainties decrease discovery reach for extra D's at LHC from 5 TeV →2 Tev [Ferrag]

Inclusive jet cross sections at LHC with PDF uncertainties (for fixed rapidity intervals)



Stump, Huston…, hep-ph/030313

Need Monte Carlos which include NLO properly

- Match NLO with parton shower
- Subtract terms which are included in parton shower from NLO result
  - At high p_T, NLO
  - At low p_T, MC

MC@NLO

$pp \rightarrow t\bar{t}$ at LHC



Frixione, Nason & Webber, hep-ph/0305252

The bottom line:

To test models of new physics we need:

Precision calculations

Higher order Monte Carlos

Much hard work remaining


Further motivation for TeV
Scale from Astrophysics…

The ASTRO/particle Connection

- 23% of universe is cold dark matter:

$$\Omega_{CDM}h^2 = .1126\ ^{+.0161}_{-.0181}$$

- WMAP (and others): Cosmology is a precision science $\longrightarrow$ Implications for particle physics!



Connection between particle/cosmology



Cold dark matter points to TeV Scale

- SUSY models have natural dark matter candidate
- LSP is neutral and weakly interacting
  - Assuming R parity
- On general grounds, LSP contributes correct relic density if mass is 300 GeV-1 TeV
- WMAP results imply M_χ <500 GeV for tan β < 40 in mSugra type models
- SUSY particles within reach of LHC and LC

Ellis, Olive, Santoso, & Spanos, hep-ph/0303043

mSugra limits from CDM



WMAP CDM



Excluded
by b→sγ

$M_\tau < M\chi$

Ellis, Olive, Santoso, Spanos, hep-ph/0303043

Extra dimension models have natural dark matter candidate

- Extra dimensions give tower of new Kaluza-Klein states
- Universal extra dimensions: all SM fields propagate in extra dimensions [Appelquist, Cheng, Dobrescu, hep-ph/0012100]
- Momentum conservation at vertices in extra D gives conserved KK number
- Lightest particle stable, can be neutral

KK WIMP



KK Wimps have TeV Scale





KK Wimps not Majorana→

efficient s-channel annihilation

Correct relic density with higher mass than LSP

Servant & Tait, hep-ph/0206071

Constructive interference of different contributions

→Lower bound on σ

Cheng, Feng, Matchev, hep-ph/0207125

The Importance of Spin

- Universal extra dimension models have new particles with same coupling and spin as SM particles
- SUSY models have new particles with same couplings, but different spin
- Spin is important tool to help distinguish between models

Spin is straightforward at Linear Collider

Threshold scan of $e^+e^- \rightarrow ZH$



Miller, hep/ph/0102023

Outlook

- Physics at TeV scale will be rich!
  - *Lots of new ideas*
  - *Connection with astrophysics*
- LHC/LC will probe TeV scale
  - *Tremendous progress in our understanding*
- Surprises await us!



Crystal Ball