



**Università
di Genova**



Jet Flavour summary

Les Houches 2023

Simone Marzani & Andreas Hinzmann







**This isn't working,
maybe we should just try
ice-cream therapy**



Yeah, but what **flavour**???


This isn't working,
maybe we should just try
ice-cream therapy

Soft Drop Flavour (SDF)



- Safe at NNLO with Jade reclustering (but not beyond)
- Only looks at information in the original anti- k_t jet
- SD only used for flavour determination, no change in the jet kinematics

Flavour anti- k_t (CMP)



Proposed modification:

A soft term designed to modify the distance of flavoured pairs.

$$d_{ij}^{(F)} = d_{ij} \begin{cases} \mathcal{S}_{ij} & i,j \text{ is flavoured pair} \\ 1 & \text{else} \end{cases}$$

$$\mathcal{S}_{ij} \equiv 1 - \theta \left(1 - \kappa_{ij}\right) \cos\left(\frac{\pi}{2} \kappa_{ij}\right) \quad \text{with} \quad \kappa_{ij} \equiv \frac{1}{a} \frac{k_{T,i}^2 + k_{T,j}^2}{2k_{T,\max}^2}.$$


- New jet algorithm to cluster together flavoured pairs first
- Small deviations from anti- k_t kinematics
- IRC sensitivity beyond NNLO can be fixed using a modified metric*

$$\mathcal{S}_{ij} \rightarrow \bar{\mathcal{S}}_{ij} = \mathcal{S}_{ij} \frac{\Omega_{ij}^2}{\Delta R_{ij}^2}$$

$$\Omega_{ik}^2 \equiv 2 \left[\frac{1}{\omega^2} (\cosh(\omega \Delta y_{ik}) - 1) - (\cos \Delta \phi_{ik} - 1) \right]$$

* see [2306.07314]

Flavour Dressing (GHS)



- start with a set of flavour agnostic jets $\{j_k\}$
- define flavoured clusters in the event $\{f_i\}$
- associate flavour clusters to jet (association criterion)
- determine the total flavour of a jet (accumulation criterion):

- algorithm only used to assign a flavour label, no kinematic modifications
- potential IRC sensitivity with many hard partons and 2 soft emissions*

* see [2306.07314]

Gauld, Huss, Stagnitto [2208.11138]





FIG. 20. Example $\mathcal{O}(\alpha_s^2)$ configuration that yields an issue for the GHS algorithm. There are four hard particles (that one can imagine recoiling against a hard gluon or electroweak system on the other side of the event), a collinear emission of a hard gluon g from one of the flavoured particles (the q), which then splits collinearly to a flavoured pair $q'\bar{q}'$.

e.g. $Zb\bar{b}$ @ N4LO

Interleaved Flavour Neutralisation (IFN)





Flavours at Les Houches

- A common framework for the 4 algorithms was created and made public <https://github.com/jetflav>
 - First stepping stone towards detailed comparisons
1. Comparisons in NNLO calculations (at parton level)
 2. Comparison with LO + parton shower and, eventually, NLO + parton shower, with and without hadronisation corrections
 3. Training machine-learning based b-taggers on jets with different b-labels and compare performances with standard anti- k_t



NNLO studies: Z+b jet (R=0.5)



interesting shape difference at low p_T : it deserves further investigation!



- all new algorithms are IRC safe at this order (anti-kt is not!) and in reasonable agreement
- CMP and SDF behaves very similarly but they differ wrt GHS/IFN by a few percent

amazing work by Rene!

NNLO studies: WH(bb)

massless b , NNLO

NNLO







- with both massless and massive b quarks

thanks to Arnd and Raoul!

LO-PS (Pythia parton level)




- CMS 8 TeV analysis but identifying **strange quarks** rather than b's to highlight differences



- For inclusive η, p_T distributions only CMP shows a tiny deviation from anti- k_t kinematics (baseline)
- For η_s, p_{T_s} distributions IRC sensitivity of SDF at higher order becomes visible, while the other 3 algorithms behave similarly
- It would be interesting to see hadronisation corrections to this picture

thanks Silvia! ... even if it's not your favourite shower...


LO-PS (Pythia hadron level)



- first hadron-level studies show larger differences... something to investigate further

thanks Federico!

LO-PS (Herwig parton & hadron level)



- First step to understand the behaviour of these algorithms on hadrons
- longer runs are needed to draw conclusions
- it would be interesting to also see correlations plots

thanks to Andrzej & the Herwig team!

Lessons learnt and future plans

- First fixed-order studies here at Les Houches show are reassuring: the algorithms behave as expected
 - When we add parton shower and hadronisation the picture changes but we haven't reached a full understanding yet
 - It's important to make these comparison more detailed and consistent (all stages of the simulation for each MC generator)
 - This will allows us to understand better the use of b-labelled jets as input for ML training and for data/theory comparisons
 - We don't always want the same thing (e.g. the discussion we had about double b-tagging) but it's important to record any useful information
-
- We have an overleaf document
<https://www.overleaf.com/project/648ab3e1c164ede47c68c368>
 - We will setup a mailing list for those of you interested in follow-up studies

Lessons learnt and future plans

- First fixed-order studies here at Les Houches show are reassuring: the algorithms behave as expected
 - When we add parton shower and hadronisation the picture changes but we haven't reached a full understanding yet
 - It's important to make these comparison more detailed and consistent (all stages of the simulation for each MC generator)
 - This will allows us to understand better the use of b-labelled jets as input for ML training and for data/theory comparisons
 - We don't always want the same thing (e.g. the discussion we had about double b-tagging) but it's important to record any useful information
-
- We have an overleaf document <https://www.overleaf.com/project/648ab3e1c164ede47c68c368>
 - We will setup a mailing list for those of you interested in follow-up studies



Thanks everyone for the great work!